

Growth Performance of *Clarias gariepinus* (Burchell, 1822) Fed With *Vigna subterranea* at Different Inclusion Levels

Aminu Saidu ^{*1} , Amina Garba Babayi ² , Basira Ibrahim ²

1.Department of Applied Biology, Kaduna Polytechnic, Kaduna, Nigeria.

2.Department of Biological Sciences, Kaduna State University, Kaduna, Nigeria.

Abstract

Vigna subterranea is an herbaceous legume from the family Fabaceae reported to contain 14-24% crude protein. The overdependence on fishmeal has led to inflation in the price of the fish feed thereby making fishes expensive. The experiment aims at evaluating the growth performance of *Clarias gariepinus* fed with *Vigna subterranea* at different inclusion levels. The effect of *Vigna subterranea* (Bambara nut) as an ingredient in the diets of *Clarias gariepinus* with an average body weight of 5.97 ± 15 was evaluated over eight (8) weeks feeding trial. Six experimental diets were formulated at 0%, 10%, 20%, 30%, 40% and 50% inclusions of Bambara nut meal (BNM) and a control (commercial feed) labelled T1, T2, T3, T4, T5, T6 and T7 respectively. The seeds were processed by soaking in water, air dried, toasted and ground using mortar pestle to obtain Bambara nut meal used to replace fishmeal at different inclusion levels. The experiment was conducted using 25liter capacity rubber tanks at the stocking density of 10 fingerling per tank and replicated three times. Fish fed T6 (50% inclusion of BNM) produced the best result, it was followed by T5 and T4 though there was no significant difference between the control (0% and commercial feed) and the formulated diets at p-value (<0.05). The study shows that processed BNM can replace fish meal in the diet of African Catfish at graded levels of up to 50%.

Keywords: *Fishmeal, Vigna subterranea, Inclusion and Clarias gariepinus.*

INTRODUCTION

Vigna subterranea also called Bambara-nut is one of the well-known plant in Africa, is been favored in terms of nutritional value and tolerance to adverse environmental conditions. It has been recorded as the third most important legumes plant after ground nuts and cowpea respectively. It is a leguminous plant common to the groundnut which is usually grown for its underground seeds especially in the middle belt region and Enugu state of Nigeria [13 , 20]. The seeds are often eaten raw when immature while the hard-mature seeds have to be roasted or boiled for consumption. The crude protein level of the seed ranges from 14-24% protein, about 60% carbohydrate, and it is higher in essential amino acids like isoleucine, leucine, lycine, phenylalanine, threonine, valine and methionine than most other grain legumes including groundnut. It contains 6-12 % oil an amount that is lesser than that in peanuts [14].

Bambara nut has not been effectively cultivated or underutilized especially in hostile tropical environments. The utilization of Bambara nut is also limited by the presence of growth inhibitors such as trypsin and chymotrypsin inhibitors, phytates, nitrates and cyanogen' s. Traditional processing techniques such as soaking, pouting or cooking have limited effects on the improvement of protein quality of grain legumes and deficiencies in some essential amino acids. In view of challenges posed by fishmeal in feed industry, research into alternative protein source such as Bambara nut in the diet of *Clarias gariepinus* becomes imperative. The essential amino acid content of Bambara groundnut such as lysine 6.82g/16gN, methionine 1.85g/16gN and cysteine 1.24g/16gN is comparable to that of soybean with 6.24g/16gN lysine, 1.14g/16gN methionine and 1.80g/16gN cysteine [6].

Processing methods such as dehulling, milling, soaking, cooking, fermentation, autoclaving, roasting and frying have been found to reduce/ eliminate anti-nutritional factors present in feed ingredients. Soaking, drying and milling processes significantly reduced the trypsin inhibitor, tannins, phytate and hemagglutinin of the Bambara nut. The nutritional implication of these reductions in the concentration of trypsin inhibitor is that it will lead to improvement in protein digestibility. Fermentation is a simple and cheap method to decrease the anti-nutritional factors contained in plants by-products [17].

The aim of this research was to evaluate the growth performance of Catfish fed processed *Vigna subterranea* for eight weeks feeding trials.

MATERIALS AND METHODS

Study Area

The research was carried out in the Zoology Garden, Department of Biological Sciences, Kaduna State University located at Latitude 10.31⁰ North and Longitude 7.26⁰ East and 6.14 meter above the Sea Level

Source of *Vigna subterranea*

Twenty five kilogram (25Kg) of *Vigna subterranea* were sourced from Station market within Kaduna metropolis, and taken to Biological Sciences Department, Kadunana State University for identification

and authentication. The specimen was assigned the reference number kasu/bsh/. This method was adopted by [4].

Source of Experimental fish

One hundred and eighty (180) mixed sexes fingerlings of *Clarias gariepinus* were procured from Federal Ministry of Agricultural and Rural Development Live House Mando Road Kaduna State. The fish were harvested, kept overnight and transported early in a 25liter capacity jerry can to the experimental site. The fish were fed with 2mm sized commercial feed containing CP of 40% for two weeks acclimatization period [7].

Preparation and Processing of *Vigna subterranea*

Vigna subterranea seeds were washed and soaked in water for twenty-four hours, air dried and pulverized into fine powder using mortar and pestle to obtain Bambara nut meal tagged (BNM). This method was adopted by [18].

Feed formulation of the experimental diet

The experimental feeds were formulated using Pearson's Square Method, incorporating maize as an energy source, groundnut cake and fishmeal as protein sources, and *Bambara nut* meal as a partial fishmeal replacement at inclusion levels of 0%, 10%, 20% 30%, 40% and 50%. Other ingredients, including vitamin premix, wheat offal, rice bran, methionine, lysine, and water, were also added. The mixture was pelleted to a size of 3 mm using a local pelleting machine at Uyama Agro Nigeria Limited, Kaduna. This method was adopted by [5].

Experimental Design

The experiment consists of seven treatments each representing 0, 10, 20, 30, 40, and 50 percent inclusions of Bambara nut meal to replace fishmeal in diet T1, T2 T3, T4, T5 and T6 that were formulated. Each of these treatments were replicated thrice. The controls (T7 or commercial feed) have no inclusion of Bambara nut meal. 21 rubber tanks of 25liters capacity were used at the stocking density of 10 fish per tank. Feeding was twice a day at 5% body weight in the morning around 9:00am, and evening around 4:00pm throughout the experimental period. Water was changed every 24hours in order to avoid stressing the fish throughout the eight weeks experimental period as adopted by [2].

Growth performance

Weight and length of the experimental fish were taken using weighing balance and meter rule respectively once in every two weeks to determine the total weight gain (TWG) and total length gain as adopted by [4].

Performance evaluation

During the experiment, fish performance was based on productivity indices on growth performance and nutrient utilization efficiencies as described by [1].

Total feed intake (TFI)

Total feed intake was estimated by summing the weekly feed intakes during the period of the experiment [19].

Total weight gain (TWG)

Total weight gain was obtained as the difference between the initial weight and the final weight gained of the experimental fish using the formula below:

Total weight gain (TWG) = final weight – initial weight [3].

Total percentage weight gain (TPWG %)

Total percentage weight gain was calculated using the formula below:

TPWG = Total weight gained / Initial weight x 100% [3].

Feed conversion ratio

From the feed consumed by each group of fish and weight gained, the feed conversion ratio (FCR) was calculated using the expression adopted by

FCR = Feed Intake / Net weight gain [8].

Protein Efficiency Ratio

Protein Efficiency Ratio (PER) was calculated from the weight gain obtained from the test subject divided by its intake of a particular feed protein during the test period. Hence

PER = Gain in body mass (g)/ Protein intake (g) [10].

Data Analysis

Analysis of Variance (ANOVA) was used to evaluate significant differences among experimental fish fed different inclusions of *Vigna subterranea* and the controls. P-value < 0.05 was considered to be significant. Statistical package used was Statistical Analysis System (SAS) version 9.4.

RESULTS**Growth performance of *Claris gaiepinus* fed with *Vigna subterranea* for Eight Weeks Feeding Trials****Initial Length of *Claris gaiepinus* Fed with *Vigna subterranea* for Eight Weeks Feeding Trials**

The average initial weight of *Clarias gariepinus* at the commencement of the experiment ranged from 5.97 ± 0.15 to 6.10 ± 0.44 g. Statistical analysis ($p > 0.05$) revealed no significant difference in the initial weights across all treatments, indicating a uniform distribution of the experimental fish (Table 1).

Final Length gain of *Clarias gariepinus* Fed with *Vigna subterranea* for Eight Weeks Feeding Trials

Table 1 shows that the final length gain of *Clarias gariepinus* fed Bambara nut meal at different inclusion levels for eight weeks feeding trials was between 15.1 ± 0.06 to 16.2 ± 0.06 . At p -value > 0.05 , the least significant difference (LSD) indicated that there was significant difference between the treatments and the control fed commercial feed.

Table 1 Average Body Length (cm) of the *Clarias gariepinus* fed *Vigna subterranea* for eight weeks feeding trials.

Parameter	T1 (0%)	T2 (10%)	T3 (20%)	T4 (30%)	T5 (40%)	T6 (50%)	commercial
W1	5.80 ± 0.25^e	6.03 ± 0.25^e	6.10 ± 0.36^e	5.97 ± 0.15^e	6.10 ± 0.36^e	6.10 ± 0.44^e	6.10 ± 0.26^e
W2	10.6 ± 0.15^d	10.7 ± 0.12^d	10.8 ± 0.20^d	10.9 ± 0.12^d	10.8 ± 0.20^d	10.8 ± 0.20^d	10.9 ± 0.10^d
W4	12.1 ± 0.10^c	12.2 ± 0.06^c	12.5 ± 0.06^c	12.4 ± 0.12^c	12.1 ± 0.06^c	12.1 ± 0.06^c	12.1 ± 0.06^c
W6	14.6 ± 0.21^b	14.6 ± 0.06^b	14.7 ± 0.06^b	14.8 ± 0.00^b	14.6 ± 0.00^b	14.6 ± 0.06^b	14.6 ± 0.06^b
W8	15.8 ± 0.12^a	15.8 ± 0.06^a	16.2 ± 0.06^a	16.0 ± 0.06^a	15.1 ± 0.06^a	15.1 ± 0.06^a	15.1 ± 0.06^a
p value	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001

Values are given as mean standard deviation. In each column, means with the same letter are not significantly different ($p < 0.05$), T1= Treatment 1, T2= Treatment 2, T3= Treatment 3, T4= Treatment 4, T5= Treatment 5, T6= Treatment 6

Average Initial Weight of *Clarias gariepinus* Fed with *Vigna subterranea* for Eight Weeks Feeding Trials

The average initial weight of the experimental fish before the start of the feeding with Bambara nut meal for eight weeks was between 4.33 ± 0.05 and 4.83 ± 0.10 . The result of Analysis of Variance revealed that there was no significant difference at p -value (< 0.05) (Table 2).

Final Weight Gain of *Clarias gariepinus* Fed with *Vigna subterranea* for Eight Weeks Feeding Trials

The final weight of the experimental fish fed Bambara nut meal for eight weeks was between 25.5 ± 0.06^a and 22.7 ± 0.15^a . The highest result was found in Treatment 1 and 7 that produced 22.7 ± 0.15^a and 26.7 ± 0.10^a respectively. This is followed by Treatment 2, 4, 5, 6 and 7 with the values of 22.66 ± 0.10^a , 25.5 ± 0.06^a , 22.6 ± 0.10^a , 22.6 ± 0.10^a and 22.6 ± 0.6^a respectively. The result of analysis of variance (ANOVA) shows that there was no significant difference at p-value (<0.05) with the control (Table 2).

Table 2 Mean Body weight (g) of the *Clarias gariepinus* fed Bambara nut meal for eight weeks Feeding Trials

Parameter	T1 (0%)	T2 (10%)	T3 (20%)	T4 (30%)	T5 (40%)	T6 (50%)	Commercial
W1	4.50 ± 0.17^e	4.33 ± 0.05^e	4.63 ± 0.06^e	4.50 ± 0.11^e	4.53 ± 0.12^e	4.63 ± 0.21^e	4.83 ± 0.10^e
W2	6.67 ± 0.12^d	6.63 ± 0.15^d	6.63 ± 0.12^d	6.70 ± 0.17^d	6.80 ± 0.17^d	6.73 ± 0.12^d	6.67 ± 0.06^d
W4	10.7 ± 0.12^c	11.1 ± 0.06^c	13.5 ± 0.06^c	12.3 ± 0.06^c	10.5 ± 0.06^c	10.9 ± 0.06^c	10.8 ± 0.06^c
W6	15.2 ± 0.10^b	15.5 ± 0.10^b	18.4 ± 0.10^b	17.5 ± 0.10^b	15.4 ± 0.10^b	15.6 ± 0.06^b	15.7 ± 0.06^b
W8	22.7 ± 0.15^a	22.66 ± 0.10^a	26.7 ± 0.10^a	25.5 ± 0.06^a	22.6 ± 0.10^a	22.6 ± 0.10^a	22.6 ± 0.6^a
p value	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001

Values are given as mean standard deviation. In each column, means with the same letter are not significantly different ($p < 0.05$), T1= Treatment 1, T2= Treatment 2, T3= Treatment 3, T4= Treatment 4, T5= Treatment 5, T6= Treatment 6

Specific Growth Rate (SGR) of *Clarias gariepinus* Fed with *Vigna subterranea* for Eight Weeks Feeding Trials

The specific growth rate (SGR) of the experimental fish fed Bambara nut meal for eight weeks ranged from 2.77 ± 0.03 - 3.13 ± 0.00 . Treatment 3 have the highest value of 3.13 ± 0.00 , followed by Treatment 2 with 2.91 ± 0.02 . The least value was observed in the control (commercial feed) with 2.77 ± 0.03 . The result of analysis of variance revealed that there was significant difference at p-value (<0.05) when compared with the control (Table 3).

Feed Conversion Ratio (FCR) of *Clarias gariepinus* Fed with *Vigna subterranea* for Eight Weeks Feeding Trials

The feed conversion ratio of the experimental feed fed Bambara nut meal for eight weeks ranged from 92.9-113.4. The best result was observed in Treatment 5 with 113.4, followed by Treatment 1 with 113.2. the least value was observed in Treatment 3 with 92.9 (Table 3).

Survival Rate of *Clarias gariepinus* Fed with *Vigna subterranea* for Eight Weeks Feeding Trials

The result of analysis of variance shows that there was Hundred percent survival rate of the experimental fish fed experimental feed (Bambara nut meal) at different inclusion level (Table 3)

Table 3 Feed Utilization Parameters Survival Rate of *Clarias gariepinus* Fed with *Vigna subterranea* for Eight Weeks Feeding Trials

Parameter	T1 (0%)	T2 (10%)	T3 (20%)	T4 (30%)	T5 (40%)	T6 (50%)	commercial
FBG (g)	22.7±0.15 ^c	22.6±0.10 ^c	26.7±0.10 ^a	25.5±0.06 ^b	22.6±0.10 ^c	22.6±0.10 ^c	22.6±0.6 ^c
IBW (g)	4.50±0.17 ^d	4.33±0.05 ^e	4.63±0.06 ^b	4.50±0.11 ^d	4.53±0.12 ^c	4.63±0.21 ^b	4.83±0.10 ^a
WG (g)	18.2±0.31 ^c	18.2±0.06 ^c	22.1±0.15 ^a	21.0±0.06 ^b	18.1±0.06 ^{cd}	17.9±0.21 ^{cd}	17.8±0.10 ^d
PWG (%)	404.3±22.6 ^b	409.8±4.79 ^b	476.3±9.05 ^a	467.4±1.28 ^a	398.7±11.0 ^b	388.4±22.0 ^{bc}	370.9±8.91 ^c
SGR (%)	2.89±0.08 ^b	2.91±0.02 ^b	3.13±0.00 ^a	2.87±0.04 ^b	2.87±0.04 ^b	2.83±0.08 ^{cb}	2.77±0.03 ^c
FI (g)	2059.4	2056.5	2054.6	2076.5	2052.9	2053.8	2068.5
FCR	113.2	112.9	92.9	98.9	113.4	114.7	116.2
PI	21.4	22.2	22.3	21.4	21.1	20.9	20.7
PER	0.85	0.82	0.98	0.97	0.86	0.86	0.86
Survival (%)	100	100	100	100	100	100	100

Values are given as mean standard deviation. In each column, means with the same letter are not significantly different ($p < 0.05$). FBW: Final Body Weight; IBW: Initial Body Weight; WG: weight gain; PWG: percentage weight gain; SGR: specific growth rate; FI: feed intake; FCR: feed conversion ratio; PER: protein efficiency ratio $FCR = \text{Feed Intake (FI)} / \text{Weight Gain (WG)}$ Protein Intake (PI) = FI * percent protein in feed $PER = WG / PI$

DISCUSSION

The growth performance of the *Clarias gariepinus* fed with *Vigna subterranea* at different inclusion levels revealed that, the average initial weight at the onset of the feeding trials, there was no significant difference among the Treatments and the control at p-value (<0.05). This also showed that all the treatments and the control were given an equal chance to compete in attaining a particular weight gain. While the average weight gain at the end of eight weeks feeding trials is in conformity with result

obtained by [12] who worked on Evaluation of the partial replacement of dietary fish meal with fermented or untreated soybean meal in juvenile silver barb, *Barbonymus gonionotus*. This is in conformity with work of [15], who experiment on growth performance of *Oreochromis niloticus* fed Bambara nut and concluded that incorporating up 50% Bambara nut meal does not have detrimental effect on the growth. This is consistent with the result obtained by [11] Assessment of growth performance of African catfish (*Clarias gariepinus*) fed with feed produced from blend of pigeon pea (Cajanuscajan) Bambara groundnut (*Vigna subterranea*) and fish meal and concluded that it is possible to produce quality Catfish feed from blend Pigeon pea, Bambara nut and fishmeal.

However, this result is similar with that of [9], who worked on Replacement effects of soybean meal with sesame seed cake on growth, biochemical body composition, and economic efficiency of *Cyprinus carpio* formulated diet and concluded that comparable growth performance and measured physiological responses indicated that Soy bean meal could be replaced by 75 to 100% Sesame seed cake in the formulated diet of *Cyprinus carpio* juveniles. The result also agreed with that of [16] who worked on Replacing fish meal and fish oil in industrial fish feeds. The result is also consistent with that of [3], who worked on influence of dietary protein content on growth performance, feed efficiency, condition factor, and length-weight relationship in *Cyprinus carpio* during the summer season.

The result of feed conversion ratio obtained in this research is similar with that of [21], who worked on effect of replacement of Soyabean (glycine max) meal with Bambara nut (*Vigna subterranea*) hull meal in the diet of African Catfish (*Clarias gariepinus*) juveniles, they concluded that Bambara nut hull meal contains necessary growth factors required for *Clarias gariepinus* juvenile and 25% is recommended for use.

The survival rate of *Clarias gariepinus* fingerlings fed *Vigna subterranea* suggested that all diets including the control were suitable for the fish and met its nutritional requirement which may be due to the processing method applied to the seeds. This result is in conformity with that of [21], who acknowledged that *Clarias gariepinus* can tolerate and survive more with the diet containing Bambara nut meal than that of Soybean meal.

CONCLUSIONS

The study reveals that the *Clarias gariepinus* fed Bambara nut meal at different inclusion levels and commercial (control) produced a final weight gain that ranges between 22.6 ± 0.10 and 26.7 ± 0.10 . It is concluded that Bambara nut Meal can replace fish meal up 50% inclusion level.

REFERENCES

- [1] Abo-Taleb, H. A., El-Feky, M. M., Azab, A. M., Mabrouk, M. M., Elokaby, M. A., Ashour, M., Mansour, A. T., Abdelzaher, O. F., Abualnaja, K. M., & Sallam, A. E. (2021). Growth performance, feed utilization, gut integrity, and economic revenue of grey mullet, *Mugil*

- cephalus, fed an increasing level of dried zooplankton biomass meal as fishmeal substitutions. *Fishes*, 6(3), 38.
- [2] Afia, O. E., David, G. S., & Umoren, I. U. (2020). Varying feeding levels and the growth response of North African catfish. *South Asian Res. J. Agric. Fish*, 2, 56-62.
- [3] Al Sulivany, B. S., Hassan, N. E., & Mhammad, H. A. (2024). Influence of Dietary Protein Content on Growth Performance, Feed Efficiency, Condition Factor, and Length-Weight Relationship in *Cyprinus carpio* during the Summer Season. *Egyptian Journal of Aquatic Biology & Fisheries*, 28(2).
- [4] Aminu, S., & Ibrahim, B. (2022). Growth Performance of *Clarias gariepinus* (Burchell, 1822) Fed Processed *Moringa oleifera* (LAM.) Seed Meal. *International Journal of Science for Global Sustainability*, 8(2), 7-7.
- [5] Apollo, M. A., Ogello, E., Balirwa, J. S., & Bassa, S. (2021). Dietary Fishmeal Substitution by Peanut-Based Meals in Nile Tilapia (*Oreochromis Niloticus* L.): Effect of Pond Water Quality on Biomass Production. *Uganda Journal of Agricultural Sciences*, 21(1), 57-67.
- [6] Bala, K., & Rano, M. (2022). Influence of processing method on the proximate composition and anti-nutrient content of bambara nut (*Vigna subterranea* L.). *Nigerian Journal of Animal Science and Technology (NJAST)*, 5(2), 92-98.
- [7] Cui, X., Huang, X., Chen, X., Li, H., Wu, Y., Yang, Z., Liu, Z., Feng, R., Xu, J., & Wei, C. (2024). Influence of Starvation on Biochemical, Physiological, Morphological, and Transcriptional Responses Associated with Glucose and Lipid Metabolism in the Liver of Javelin Goby (*Synechogobius hasta*). *Animals*, 14(18), 2734.
- [8] Elvy, J. E., Symonds, J. E., Hilton, Z., Walker, S. P., Tremblay, L. A., Casanovas, P., & Herbert, N. A. (2022). The relationship of feed intake, growth, nutrient retention, and oxygen consumption to feed conversion ratio of farmed saltwater Chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture*, 554, 738184.
- [9] Hekmatpour, F., Amiri, F., Yooneszadeh Fashalami, M., Nazemroaya, S., Sadr, A., Mousavi, S., Mortezaavizadeh, S., Kianersi, F., Ahangarzadeh, M., & Houshmand, H. (2023). Replacement effects of soybean meal with sesame seed cake on growth, biochemical body composition, and economic efficiency of *Cyprinus carpio* formulated diet. *Iranian Journal of Fisheries Sciences*, 22(3), 678-700.
- [10] Hoskin, J. M. (2023). Improved Performance of the Reference Casein Control Group in AOAC Official Method SM 960.48 (Protein Efficiency Ratio, Rat Bioassay) for Determining the Biological Quality of Protein in Infant Formula. *Journal of AOAC International*, 106(1), 1-5.
- [11] Ishiwu, C., Opara, A., Obiegbuna, J., & Okeke, P. (2020). Assessment of growth performance of African catfish (*Clarias gariepinus*) fed with feed produced from blend of pigeon pea (*Cajanus cajan*) bambara groundnut (*Vigna subterranea*) and fish meal. *Global Journal of Agricultural Sciences*, 19(1), 17-26.
- [12] Jahan, H., Tumpa, I. J., Qasem, W. A., Moniruzzaman, M., Pervin, M. A., Akter, R., Omri, A., Min, T., & Hossain, Z. (2021). Evaluation of the partial replacement of dietary fish meal with

- fermented or untreated soybean meal in juvenile silver barb, *Barbonymus gonionotus*. *Frontiers in nutrition*, 8, 733402.
- [13] Jideani, V. A., & Jideani, A. I. (2021). *Bambara groundnut: Utilization and future prospects*. Springer.
- [14] Mbuma, N. W., Labuschagne, M., Siwale, J., & Hugo, A. (2022). Diversity in seed protein content, selected minerals, oil content and fatty acid composition of the Southern African Bambara groundnut germplasm collection. *Journal of Food Composition and Analysis*, 109, 104477.
- [15] Mohammed, M., Absalom, K., Awolumate, S., Mopho, F., Anthony, B., Izang, A., & Oshibanjo, D. (2024). GROWTH PERFORMANCE OF TILAPIA (*OREOCHROMIS NILOTICUS*) FED BAMBARA NUT. *Nigerian Journal of Animal Production*, 1975-1979.
- [16] Oliva-Teles, A., Enes, P., Couto, A., & Peres, H. (2022). Replacing fish meal and fish oil in industrial fish feeds. *Feed and Feeding Practices in Aquaculture*, 231-268.
- [17] Pokharel, U. (2022). *EFFECT OF PROCESSING METHODS ON ANTINUTRITIONAL FACTORS PRESENT IN GREEN GRAM [MUNG BEAN]* Department of Food Technology Central Campus of Technology, Dharan Institute ...].
- [18] Saidu, A., Abdulrahman, A., & Imam, Z. (2023). Effect of processing methods on the proximate and phytochemical constituents of Moringa Oleifera (Lamarck, 1785) leaves. *Science World Journal*, 18(2), 272-275.
- [19] Sun, H., Chen, D., Cai, H., Chang, W., Wang, Z., Liu, G., Deng, X., & Chen, Z. (2022). Effects of fermenting the plant fraction of a complete feed on the growth performance, nutrient utilization, antioxidant functions, meat quality, and intestinal microbiota of broilers. *Animals*, 12(20), 2870.
- [20] Wanyama, A. W. (2018). *Evaluation of Phytoconstituents, Antioxidants Potential, Cytotoxic, Antimicrobial Activities and Mineral Composition of Vigna subterranea (L) Verdic. Extracts JKUAT-COHES*].
- [21] Yusuf, A., Yunisa, A., Buba, W., Anayeokwu, S., & Emmanuel, V. (2024). EFFECT OF REPLACEMENT OF SOYABEAN (*Glycine max*) MEAL WITH BAMBARA NUT (*Vigna subterranea*) HULL MEAL IN THE DIET OF AFRICAN CATFISH (*Clarias gariepinus*) JUVENILES. *International Journal of Global Affairs, Research and Development*, 2(1), 136-145.